

Decreasing Inappropriate Care in California in Partnership with Choosing Wisely® - Summary of Measures, Data, and Targets

Sutter Health: Measures and Targets			
Target treatment/ test	<i>Avoidance of antibiotics for adults with acute bronchitis</i>	<i>Imaging for uncomplicated headache</i>	<i>Don't perform repetitive CBC and chemistry testing</i>
Measure	The percentage of adults 18–64 years of age with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription. Specification: IHA P4P Measure (HEDIS) HIGHER IS BETTER	Percent of patients with an office visit for a diagnosis of headache who received imaging for headache within 30 days. Specification: Washington State Choosing Wisely Task Force LOWER IS BETTER	Percent of hospitalized patients with orders for CBC and metabolic panel for whom the lab tests were performed as part of a “repeating order.” Specification: Sutter Health Variation Reduction Program LOWER IS BETTER
Exclusions	Included in the HEDIS specification	None	None
Codes (CPT or ICD)		Denominator: Count of patients with an office visit with a Primary Care Practitioner (Epic encounter type = office visit) with a diagnosis code for headache (ICD9=784.0) Numerator: Count of patients with an imaging order (Epic proc code= RADXXXX) within 30 days of the visit with an associated diagnosis of headache (ICD9=784.0)	Denominator: Count of CBC and Metabolic Panel labs ordered by hospitalists (Epic proc code = LABCBCA, LABBMPG, LABCMPG, LABCBCND) Numerator: Count of labs that were ordered as repeating (repeating = more than one Epic “child” order)
Baseline Data	46% ¹	9% ¹	49% ²
Utilization Period 1	45% ³	9% ³	37% ⁴
Target	55.2%	7.2%	39.2%
Data Source	EHR	EHR	EHR

¹ Baseline data covers 3/14-2/15

² Baseline data covers 9/13-8/14

³ Utilization Period 1 covers 3/15-9/15

⁴ Utilization Period 1 covers 9/14-9/15

Sharp Rees-Stealy Medical Group: Measures and Targets

Target treatment/ test	<i>Avoidance of antibiotics for adults with acute bronchitis</i>	<i>Imaging for nonspecific low back pain</i>	<i>Preoperative stress testing</i>
Measure	The percentage of adults 18–64 years of age with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription. Specification: IHA P4P Measure (HEDIS) HIGHER IS BETTER	The percentage of members with a primary diagnosis of low back pain who did not have an imaging study (X-ray, MRI, CT scan) within 28 days of the diagnosis. Specification: IHA P4P Measure (HEDIS) HIGHER IS BETTER	Use of preoperative stress testing in patients scheduled to undergo low- and/or intermediate-risk non-cardiac surgery. Report as a rate per thousand. Specification: Internal, with guidance from NQF specification #0672 LOWER IS BETTER
Exclusions	Included in the HEDIS specification	HEDIS specification/codes with no modifications.	None
Codes (CPT or ICD)			<ul style="list-style-type: none"> • CPT: 93015, 93016, 93017, 93018, 93325, 93350, 78452 • Claims Location: Outpatient Hospital (2), Doctors Office (3)
Baseline Data (CY 2014)	75.3%	83.1%	8.75 per thousand
Utilization Period 1 (Jan-Sep 2015)	73.4%	81.46%	8.42 per thousand
Target	80.2%	86.5%	7.00 per thousand
Data Source	Claims	Claims	Claims